

F/G 12/2

HEAVY TRAFFIC RESULTS FOR SINGLE SERVER QUEUES WITH DEPENDENT (--ETC(U)

APR 79 P A JACOBS

NPS55-79-009

UNCLASSIFIED

NL


|OF|

AD  
A070086

100%

END  
DATE  
FILMED

7-79  
DOC




MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

NPS55-79-009

**NAVAL POSTGRADUATE SCHOOL**  
Monterey, California



**LEVEL** *II*



**HEAVY TRAFFIC RESULTS  
FOR SINGLE SERVER QUEUES WITH  
DEPENDENT (EARMA) SERVICE  
AND INTERARRIVAL TIMES**

by

P. A. Jacobs

April 1979

Approved for public release; distribution unlimited.

79 06 20 055

AD A070086

DDC FILE COPY


NAVAL POSTGRADUATE SCHOOL  
MONTEREY, CALIFORNIA

Rear Admiral T. F. Dedman  
Superintendent

J. R. Borsting  
Provost


Reproduction of all or part of this report is authorized.


This report was prepared by:

  
P. A. Jacobs, Associate Professor  
Department of Operations Research

Reviewed by:

Released by:

  
Michael G. Sovereign, Chairman  
Department of Operations Research

  
William M. Tolles  
Dean of Research



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

| REPORT DOCUMENTATION PAGE   |                       | READ INSTRUCTIONS<br>BEFORE COMPLETING FORM                 |
|---|-----------------------|---|
| 1. REPORT NUMBER<br>14 NPS55-79-009   | 2. GOVT ACCESSION NO. | 3. RECIPIENT'S CATALOG NUMBER                               |
| 4. TITLE (and Subtitle)<br>6 Heavy Traffic Results for Single Server Queues with Dependent (EARMA) Service and Interarrival Times   |                       | 5. TYPE OF REPORT & PERIOD COVERED<br>Technical rept.       |
| 7. AUTHOR(s)<br>10 P. A. Jacobs   |                       | 6. PERFORMING ORG. REPORT NUMBER                            |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS<br>Naval Postgraduate School<br>Monterey, CA 93940  |                       | 8. CONTRACT OR GRANT NUMBER(s)<br>12 38p.                   |
| 11. CONTROLLING OFFICE NAME AND ADDRESS<br>Naval Postgraduate School<br>Monterey, CA 93940  |                       | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)   |                       | 12. REPORT DATE<br>11 April 1979                            |
|   |                       | 13. NUMBER OF PAGES<br>27                                   |
|   |                       | 15. SECURITY CLASS. (of this report)<br>Unclassified        |
|   |                       | 15a. DECLASSIFICATION/DOWNGRADING SCHEDULE                  |
| 16. DISTRIBUTION STATEMENT (of this Report)<br>Approved for public release; distribution unlimited.   |                       |   |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  |                       |   |
| 18. SUPPLEMENTARY NOTES   |                       |   |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number)<br>Cross correlated interarrival and service times<br>Multivariate exponential random variables<br>Heavy traffic<br>Weak convergence<br>EARMA sequences  |                       |   |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number)<br>Models are given for sequences of correlated exponential interarrival and service times for a single server queue. These multivariate exponential models are formed as probabilistic linear combinations of sequences of independent exponential random variables and are easy to generate on a computer. Limiting results for customer waiting time under heavy traffic conditions are obtained for these queues. Heavy traffic results are useful for analyzing the effect of correlated interarrival and service times in queues on |                       |   |

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

251 450 Gu

next page

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. ABSTRACT

*cont* such quantities as queue length and customer waiting time. They can also be used to check simulation results.



|   |  |  |  |
|---|--|--|--|
| 1. REPORT NUMBER                        |  | 2. REPORT DATE   |  |
| 3. REPORT TYPE AND DATES COVERED        |  | 4. AUTHOR  |  |
| 5. PERFORMING ORG. REPORT NUMBER        |  | 6. AUTHORING ORG. REPORT NUMBER                              |  |
| 7. CONTRACT OR GRANT NUMBER(S)          |  | 8. PROGRAM ELEMENT, PROJECT, TASK AREA AND WORK UNIT NUMBERS |  |
| 9. PERFORMING ORG. NAME(S) AND ADDRESS  |  | 10. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 11. PERFORMING ORG. NAME(S) AND ADDRESS |  | 12. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 13. PERFORMING ORG. NAME(S) AND ADDRESS |  | 14. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 15. PERFORMING ORG. NAME(S) AND ADDRESS |  | 16. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 17. PERFORMING ORG. NAME(S) AND ADDRESS |  | 18. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 19. PERFORMING ORG. NAME(S) AND ADDRESS |  | 20. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 21. PERFORMING ORG. NAME(S) AND ADDRESS |  | 22. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 23. PERFORMING ORG. NAME(S) AND ADDRESS |  | 24. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 25. PERFORMING ORG. NAME(S) AND ADDRESS |  | 26. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 27. PERFORMING ORG. NAME(S) AND ADDRESS |  | 28. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 29. PERFORMING ORG. NAME(S) AND ADDRESS |  | 30. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 31. PERFORMING ORG. NAME(S) AND ADDRESS |  | 32. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 33. PERFORMING ORG. NAME(S) AND ADDRESS |  | 34. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 35. PERFORMING ORG. NAME(S) AND ADDRESS |  | 36. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 37. PERFORMING ORG. NAME(S) AND ADDRESS |  | 38. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 39. PERFORMING ORG. NAME(S) AND ADDRESS |  | 40. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 41. PERFORMING ORG. NAME(S) AND ADDRESS |  | 42. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 43. PERFORMING ORG. NAME(S) AND ADDRESS |  | 44. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 45. PERFORMING ORG. NAME(S) AND ADDRESS |  | 46. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 47. PERFORMING ORG. NAME(S) AND ADDRESS |  | 48. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 49. PERFORMING ORG. NAME(S) AND ADDRESS |  | 50. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 51. PERFORMING ORG. NAME(S) AND ADDRESS |  | 52. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 53. PERFORMING ORG. NAME(S) AND ADDRESS |  | 54. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 55. PERFORMING ORG. NAME(S) AND ADDRESS |  | 56. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 57. PERFORMING ORG. NAME(S) AND ADDRESS |  | 58. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 59. PERFORMING ORG. NAME(S) AND ADDRESS |  | 60. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 61. PERFORMING ORG. NAME(S) AND ADDRESS |  | 62. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 63. PERFORMING ORG. NAME(S) AND ADDRESS |  | 64. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 65. PERFORMING ORG. NAME(S) AND ADDRESS |  | 66. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 67. PERFORMING ORG. NAME(S) AND ADDRESS |  | 68. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 69. PERFORMING ORG. NAME(S) AND ADDRESS |  | 70. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 71. PERFORMING ORG. NAME(S) AND ADDRESS |  | 72. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 73. PERFORMING ORG. NAME(S) AND ADDRESS |  | 74. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 75. PERFORMING ORG. NAME(S) AND ADDRESS |  | 76. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 77. PERFORMING ORG. NAME(S) AND ADDRESS |  | 78. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 79. PERFORMING ORG. NAME(S) AND ADDRESS |  | 80. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 81. PERFORMING ORG. NAME(S) AND ADDRESS |  | 82. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 83. PERFORMING ORG. NAME(S) AND ADDRESS |  | 84. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 85. PERFORMING ORG. NAME(S) AND ADDRESS |  | 86. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 87. PERFORMING ORG. NAME(S) AND ADDRESS |  | 88. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 89. PERFORMING ORG. NAME(S) AND ADDRESS |  | 90. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 91. PERFORMING ORG. NAME(S) AND ADDRESS |  | 92. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 93. PERFORMING ORG. NAME(S) AND ADDRESS |  | 94. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 95. PERFORMING ORG. NAME(S) AND ADDRESS |  | 96. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 97. PERFORMING ORG. NAME(S) AND ADDRESS |  | 98. PERFORMING ORG. NAME(S) AND ADDRESS                      |  |
| 99. PERFORMING ORG. NAME(S) AND ADDRESS |  | 100. PERFORMING ORG. NAME(S) AND ADDRESS                     |  |

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)



**HEAVY TRAFFIC RESULTS FOR SINGLE SERVER QUEUES**  
**WITH DEPENDENT (EARMA) SERVICE**  
**AND INTERARRIVAL TIMES**

by

P. A. Jacobs \*

Naval Postgraduate School  
 Monterey, California

0. ABSTRACT

Models are given for sequences of correlated exponential interarrival and service times for a single server queue. These multivariate exponential models are formed as probabilistic linear combinations of sequences of independent exponential random variables and are easy to generate on a computer. Limiting results for customer waiting time under heavy traffic conditions are obtained for these queues. Heavy traffic results are useful for analyzing the effect of correlated interarrival and service times in queues on such quantities as queue length and customer waiting time. They can also be used to check simulation results.

|                    |                                     |
|--------------------|-------------------------------------|
| Accession For      |                                     |
| NTIS GRA&I         | <input checked="" type="checkbox"/> |
| DDC TAB            | <input type="checkbox"/>            |
| Unannounced        | <input type="checkbox"/>            |
| Justification      |                                     |
| By                 |                                     |
| Distribution/      |                                     |
| Availability Codes |                                     |
| Dist               | Availand/or special                 |
| A                  |                                     |

\* Support from NSF grants ENG-77-09020 and ENG-79-01438 and ONR contract NR-42-284 is gratefully acknowledged.

## 1. INTRODUCTION

Much of the work in queueing theory deals with models in which assumptions of independent service and interarrival times are made. One way to examine the effect of correlated interarrival and service times on measures of performance in queues is to consider models in which the interarrival and service times are correlated stationary sequences of random variables having given marginal distributions and compare results for these models with those for which the interarrival and service times are independent and identically distributed with the same marginal distributions.

In [3] Jacobs and Lewis introduced a scheme for generating sequences of dependent exponential random variables. The EARMA (1,1) (exponential mixed autoregressive moving average with both autoregression and moving average of order 1) model is defined as follows. Let  $\{E_n\}$  be a sequence of independent exponential random variables with positive finite mean  $\nu^{-1}$ . Let  $\{J_n\}$  and  $\{K_n\}$  be independent sequences of independent  $\{0,1\}$ -random variables with  $P\{J_n=1\} = 1 - \beta$  and  $P\{K_n=1\} = 1 - \rho$  where  $0 \leq \beta \leq 1$  and  $0 \leq \rho < 1$  are given constants. We define

$$(1.1) \quad X_n = \beta E_n + J_n A_{n-1}, \quad n = 1, 2, \dots$$

where

$$(1.2) \quad A_n = \rho A_{n-1} + K_n E_n, \quad n = 1, 2, \dots$$

We assume that  $A_0$  is an independent exponential random variable with mean  $\nu^{-1}$ . Under this assumption  $\{X_n\}$  is a stationary sequence of dependent exponential random variables with mean  $\nu^{-1}$ .



The dependence is not Markovian in general and as was shown in [[3]

$$(1.3) \quad \text{Corr}(X_1, X_{1+n}) = \rho^{n-1} (1-\beta) [\beta(1-\rho) + (1-\beta)\rho] .$$

Note that if  $\beta = 1$  or  $\beta = 0$  and  $\rho = 0$ , then  $\{X_n\}$  is a sequence of independent exponential random variables.

The EARMA(1,1) process is formed as a probabilistic linear combination of independent random variables. This technique can be extended to form sequences of multivariate exponential random variables. This idea was briefly explored in Lewis and Shedler [7]. These models have the advantage that the marginal distribution and correlation structure of the sequence are specified separately. In section 2 we will give several ways in which to generate sequences of dependent multivariate exponential random variables from probabilistic linear combinations of independent exponentials. These stationary multivariate EARMA sequences of dependent exponentials can be used as interarrival and service times in queues. Results for measures of performance for these queues can then be compared to queues with stationary independent sequences of interarrival and service times having the same marginal (exponential) distributions. Since the correlation structure in the multivariate EARMA sequences is specified by parameters independent of the marginal distribution, any difference between results for queues with these multivariate interarrival and service sequences and the corresponding queues with independent interarrival and service sequences having the same marginal distributions will be due to the correlation structure.

Suppose that the interarrival and service times of a single server queue form stationary multivariate EARMA sequences

of dependent exponential random variables. Results of Loynes [9] give conditions for the existence of limiting distributions for the waiting time of a customer and the queue length. However, it seems to be difficult to obtain exact results even in the simplest queues with EARMA interarrival and service times (cf. Jacobs [4]).

As a result we turn our attention to the possibilities of obtaining limiting results of the heavy traffic type for queues with stationary multivariate EARMA sequences of dependent interarrival and service times. Results of this kind will indicate how dependence in queues introduced by the EARMA scheme affects quantities such as customer waiting time and queue length, at least under heavy traffic conditions. These limiting results can also be used to check simulations.

In this paper we will show that Kingman's [5] result for the equilibrium waiting time  $W$  in heavy traffic holds for queues having multivariate EARMA interarrival and service times; that is, we will show under conditions detailed in Section 3, that if the traffic intensity  $\lambda\mu^{-1}$  of the queue is close to 1, then the distribution of  $(1 - \lambda\mu^{-1})W$  is approximately exponential. The mean of this exponential distribution will in general be different from the case of independent interarrival and service times. Positive cross-correlation between the interarrival and service times will tend to decrease the mean; while positive correlation within the interarrival and service time sequences will tend to increase it.



In Section 3 we will state the heavy traffic result more precisely and present several examples of its application. Its formulation for multivariate EARMA queues is very dependent on the fact that the marginal distribution and correlation structure are specified independently of one another in EARMA formulations. In Section 4 we will give the proof of the heavy traffic result.

## 2. QUEUES WITH EARMA INTERARRIVAL AND SERVICE TIMES

We will consider a single server queue at which customers arrive and are served in the order of their arrival. The 0th customer arrives at  $t = 0$  and finds the server free. The arrival time, service time, and waiting time (excluding service) of the  $n$ th customer are denoted respectively by  $T_n = \sum_{k=1}^n X_k$ ,  $S_n$ , and  $W_n$ . In this section we will present some examples of the use of the EARMA scheme to model dependent interarrival and service times in queues.

(2.1) EXAMPLE. Let  $\{X_n\}$  be an EARMA(1,1) process with positive finite mean  $\lambda^{-1}$  and parameters  $0 \leq \beta_1 \leq 1$  and  $0 \leq \rho_1 < 1$ ; let  $\{S_n\}$  be an EARMA(1,1) process independent of  $\{X_n\}$  with positive finite mean  $\mu^{-1}$  and parameters  $0 \leq \beta_2 \leq 1$  and  $0 \leq \rho_2 < 1$ . Note that if  $\beta_1 = 1$ ,  $\beta_2 = 1$ , or  $\beta_1 = 1$ ,  $\beta_2 = 0$ ,  $\rho_2 = 0$ , or  $\beta_1 = 0$ ,  $\rho_1 = 0$ ,  $\beta_2 = 1$ , or  $\beta_1 = 0$ ,  $\rho_1 = 0$ ,  $\beta_2 = 0$ ,  $\rho_2 = 0$ , then the queue reduces to an M/M/1 queue. Other special cases include queues with Poisson arrivals and EARMA(1,1) service times independent of the arrival process, and queues with EARMA(1,1) interarrival times and independent exponential service times.

The sequence  $\{(X_n, S_n)\}$  is stationary and  $\{X_n\}$  and  $\{S_n\}$  are independent. However the  $X_n$ 's (respectively  $S_n$ 's) are positively correlated with



$$(2.2) \quad \text{Cov}(X_n, X_{n+k}) = \lambda^{-2} \rho_1^{k-1} (1-\beta_1) [\beta_1 (1-\rho_1) + (1-\beta_1) \rho_1]$$

(respectively

$$(2.3) \quad \text{Cov}(S_n, S_{n+k}) = \mu^{-2} \rho_2^{k-2} (1-\beta_2) [\beta_2 (1-\rho_2) + (1-\beta_2) \rho_2]$$

from (1.2).

Let  $\mathbb{F}_m^1$  (respectively  $\mathbb{F}_m^2$ ) be the  $\sigma$ -algebra generated by  $X_1, \dots, X_m$  (respectively  $S_1, \dots, S_m$ ) and  $\mathbb{G}_{m+k}^1$  (respectively  $\mathbb{G}_{m+k}^2$ ) be the  $\sigma$ -algebra generated by  $X_{m+k}, X_{m+k+1}, \dots$  (respectively  $S_{m+k}, S_{m+k+1}, \dots$ ). Let  $L^2(\mathbb{F}_m^i)$  (respectively  $L^2(\mathbb{G}_{m+k}^i)$ ) denote the collection of real-valued functions having finite second moment that are measurable with respect to  $\mathbb{F}_m^i$  (respectively  $\mathbb{G}_{m+k}^i$ ),  $i = 1, 2$ . If  $f_i \in L^2(\mathbb{F}_m^i)$  and  $g_i \in L^2(\mathbb{G}_{m+k}^i)$ ,  $i = 1, 2$ , then there is a  $k_0$  independent of  $f_i, g_i$ ,  $i = 1, 2$ , such that for  $k \geq k_0$

$$\begin{aligned} (2.4) \quad & |E[(f_1 f_2)(g_1 g_2)] - E[f_1 f_2] E[g_1 g_2]| \\ &= |E[f_1 g_1] E[f_2 g_2] - E[f_1] E[f_2] E[g_1] E[g_2]| \\ &= |E[f_1 g_1] \{E[f_2 g_2] - E[f_2] E[g_2]\} \\ &\quad + E[f_2] E[g_2] \{E[f_1 g_1] - E[f_1] E[g_1]\}| \end{aligned}$$

$$\begin{aligned}
(2.5) \quad & \leq E[|f_1 g_1|] 5 \rho_2^{k/2} E[f_2^2]^{1/2} E[g_2^2]^{1/2} \\
& + E[|f_2|] E[|g_2|] 5 \rho_1^{k/2} E[f_1^2]^{1/2} E[g_1^2]^{1/2} \\
& \leq 5 [\rho_1^{k/2} + \rho_2^{k/2}] E[(f_1 f_2)^2]^{1/2} E[(g_1 g_2)^2]^{1/2}
\end{aligned}$$

where (2.4) follows from the independent of  $\{X_n\}$  and  $\{S_n\}$  and (2.5) follows from (6.5) of Jacobs and Lewis [3]. Hence, if  $f$  (respectively  $g$ ) has finite second moment and is measurable with respect to the product  $\sigma$ -algebra  $\underline{F}_m^1 \times \underline{F}_m^2$  (respectively  $\underline{G}_{m+k}^1 \times \underline{G}_{m+k}^2$ ), then for  $k$  sufficiently large

$$(2.6) \quad |E[fg] - E[f] E[g]| \leq 5 E[f^2]^{1/2} E[g^2]^{1/2} [\rho_1^{k/2} + \rho_2^{k/2}].$$

Hence, the sequence  $\{(X_n, S_n)\}$  is asymptotically uncorrelated in the sense of Rosenblatt [10] and is  $\phi$ -mixing in the sense of Billingsley [1].

(2.7) EXAMPLE. In Lewis and Shedler [7] one of the schemes that is proposed for cross-correlating the arrival process and the service times is the following. Assume  $\{X_n\}$  and  $\{E_n\}$  are independent sequences of independent exponential random variables with positive finite means  $\lambda^{-1}$  and  $\mu^{-1}$  respectively. The  $n$ th interarrival time is assumed to be  $X_n$  and hence the arrival process is Poisson with rate  $\lambda$ . Let the  $n$ th service time

$$(2.8) \quad S_n = \beta E_n + J_n(\lambda \mu^{-1} B_n)$$

where

$$(2.9) \quad B_n = \rho B_{n-1} + K_n X_n$$

and  $\beta$ ,  $\rho$ ,  $\{J_n\}$  and  $\{K_n\}$  are as in (1.1) and (1.2) so that  $\{S_n\}$  and  $\{X_n\}$  are now cross-correlated sequences. We will assume that  $B_0$  has an exponential distribution with mean  $\lambda^{-1}$ .

Since  $E_n$  and  $B_n$  are independent,  $\{S_n\}$  is a stationary sequence of exponential random variables with mean  $\mu^{-1}$ . If  $\beta = 1$ , then the queue reduces to an M/M/1 queue. The  $S_n$ 's can be shown to be positively correlated with

$$(2.10) \quad \text{Cov}(S_n, S_{n+k}) = \mu^{-2} (1-\beta)^2 \rho^k, \quad k = 1, 2, \dots$$

Further, the  $n$ th service time  $S_n$  is positively correlated with the interarrival times  $X_n, X_{n-1}, \dots, X_1$  with

$$(2.11) \quad \text{Cov}(S_n, X_{n-k}) = (\lambda\mu)^{-1} (1-\beta) (1-\rho) \rho^k$$

for  $k = 0, \dots, n-1$ .

Note that  $(X_n, S_n, \dots, X_{n+k}, S_{n+k})$  is a function of  $(B_{n-1}, X_n, E_n, J_n, K_n, X_{n+1}, E_{n+1}, J_{n+1}, K_{n+1}, \dots, X_{n+k}, E_{n+k}, J_{n+k}, K_{n+k})$ . Hence, since the sequence  $\{B_n\}$  is stationary so is the sequence  $\{(X_n, S_n)\}$ . Further, by arguments similar to those used in obtaining (6.5) of Jacobs and Lewis [3], if  $f$  and  $g$  are as in (2.6), then for  $k$  sufficiently large

$$(2.12) \quad |E[fg] - E[f] E[g]| \leq 5\rho^{k/2} E[f^2]^{1/2} E[g^2]^{1/2}$$

so that the sequence  $\{(X_n, S_n)\}$  is  $\phi$ -mixing.



There are many ways to cross-correlate the interarrival and service times using the EARMA scheme beyond those given above and in Lewis and Shedler [7]. We will present one more, somewhat more general than that of Example (2.7) in which the service times were coupled back over the previous interarrival times.

(2.13) EXAMPLE. Let  $\{E_n\}$ ,  $\{I_n\}$ , and  $\{C_n\}$  be independent sequences of independent exponential random variables having respective positive finite means  $\lambda^{-1}$ ,  $\mu^{-1}$ , and 1. Let

$$(2.14) \quad X_n = \beta E_n + J_n (\lambda^{-1} A_n)$$

$$(2.15) \quad S_n = \gamma E_n + I_n (\mu^{-1} A_n)$$

with

$$(2.16) \quad A_n = \rho A_{n-1} + K_n C_n$$

where  $\beta$ ,  $\rho$ ,  $\{J_n\}$  and  $\{K_n\}$  are as in (1.1) and (1.2) and  $\{I_n\}$  is a sequence of independent binary random variables independent of everything with  $P\{I_n=1\} = 1 - P\{I_n=0\} = (1-\gamma)$  for some fixed constant  $0 \leq \gamma \leq 1$ . Again we assume that  $A_0$  is an independent random variable having an exponential distribution with mean 1. Then  $\{(E_n, E_n, A_n, I_n, J_n)\}$  is a stationary sequence of random variables and, hence, the same is true of  $\{(X_n, S_n)\}$ .



If  $\gamma = 1, \beta = 1$  or  $\gamma = 1, \beta = 0, \rho = 0$  or  $\gamma = 0, \beta = 1, \rho = 0$ , then the queue reduces to an M/M/1 queue. If  $\beta = 1, \rho = 0$  then the queue reduces to a special case of Example (2.7). Straightforward manipulation shows that in general the interarrival times  $X_n$  are positively correlated with

$$(2.17) \quad \text{Cov}(X_n, X_{n+k}) = \lambda^{-2} (1-\beta)^2 \rho^k, \quad k = 1, 2, \dots$$

The service times  $S_n$  are also positively correlated with

$$(2.18) \quad \text{Cov}(S_n, S_{n+k}) = \mu^{-2} (1-\gamma)^2 \rho^k, \quad k = 1, 2, \dots$$

The interarrival times and service times are positively cross-correlated with

$$(2.19) \quad \text{Cov}(X_n, S_{n+k}) = (1-\beta)(1-\gamma)(\lambda\mu)^{-1} \rho^k, \quad k = 0, 1, \dots;$$

and

$$(2.20) \quad \text{Cov}(X_{n+k}, S_n) = (1-\beta)(1-\gamma)(\lambda\mu)^{-1} \rho^k, \quad k = 0, 1, \dots$$

Finally, by the arguments used in obtaining (6.5) in Jacobs and Lewis [3], if  $f$  and  $g$  are as in (2.6), then for  $k$  sufficiently large

$$(2.21) \quad |E[fg] - E[f] E[g]| \leq 5\rho^{k/2} E[f^2]^{1/2} E[g^2]^{1/2}.$$

so that the sequence  $\{(X_n, S_n)\}$  is  $\phi$ -mixing.

### 3. A HEAVY TRAFFIC RESULT FOR EARMMA QUEUES

Let  $\{X_n\}$  (respectively  $\{S_n\}$ ) be the sequence of interarrival times (respectively service times) for a single server queue with first-in-first-out service discipline.

Put  $U_n = S_n - X_n$  and assume that  $\{U_n\}$  is a stationary sequence with  $E[U_n] < 0$ . Let  $W_n$  be the waiting time of the  $n$ th customer and  $W = \sup_{n \geq 0} \sum_{k=1}^n U_k$  (with the convention that  $\sum_{k=1}^0 = 0$ ). Loynes [9] has shown that if the sequence  $\{U_n\}$  is metrically transitive, then the distribution of  $W_n$  tends in the limit to the distribution function of  $W$ . Since it is usually difficult to obtain the distribution of  $W$  analytically, particularly when the service and interarrival times are dependent, we will obtain a heavy traffic limiting result for it. In this section we will give the statement of the result as well as some examples of its use leaving the proof to the next section.

For each  $\alpha > 0$  let  $Q_\alpha$  be a single server queue with FIFO queueing discipline having stationary sequences of interarrival times  $\{X_n(\alpha)\}$  with positive finite mean  $\lambda(\alpha)^{-1}$  and service times  $\{S_n(\alpha)\}$  with positive finite mean  $\mu(\alpha)^{-1}$  such that, if  $U_n(\alpha) = S_n(\alpha) - X_n(\alpha)$ , then  $\{U_n(\alpha)\}$  is a stationary sequence of random variables. Let

$$(3.1) \quad E[U_n(\alpha)] = v(\alpha) < 0; \text{Var}[U_n(\alpha)] = s^2(\alpha);$$



$$(3.2) \quad \sigma^2(\alpha) = \text{var } U_1(\alpha) + 2 \sum_{n=1}^{\infty} \text{Cov}(U_1(\alpha), U_{1+n}(\alpha));$$

$$(3.3) \quad d(\alpha) = |v(\alpha)|/\sigma(\alpha) \quad \text{and} \quad n(\alpha) = d(\alpha)^{-2}.$$

We will assume the following:

(3.4) ASSUMPTION. There exists a distribution function  $G$  so that  $\lim_{\alpha \rightarrow \infty} P\{[U_n(\alpha) - v(\alpha)]^2 \leq x\} = G(x)$  for all continuity points  $x$  of  $G$ ;  $\lim_{\alpha \rightarrow \infty} v(\alpha) = 0$ ; and  $0 < \lim_{\alpha \rightarrow \infty} \sigma^2(\alpha) = s^2 < \infty$ .

(3.5) ASSUMPTION.  $0 < \lim_{\alpha \rightarrow \infty} \sigma^2(\alpha) = \sigma^2 < \infty$ .

(3.6) ASSUMPTION. If  $f$  (respectively  $g$ ) is a function with finite second moment measurable with respect to the  $\sigma$ -algebra generated by  $U_1(\alpha), \dots, U_m(\alpha)$  (respectively  $U_{m+k}(\alpha), U_{m+k+1}(\alpha), \dots$ ), then

$$(3.7) \quad |E[fg] - E[f]E[g]| \leq \phi(k) E[f^2]^{1/2} E[g^2]^{1/2}$$

for some decreasing function  $\phi$  bounded above by 1 not depending on  $\alpha$  satisfying

$$(3.8) \quad \sum_{n=1}^{\infty} \phi(n) < \infty.$$

Assumption (3.4) concerns the convergence of the distribution of  $U_n^2(\alpha) = [S_n(\alpha) - X_n(\alpha)]^2$  and its first two moments as  $\alpha \rightarrow \infty$ . Assumption (3.6) is roughly a condition that the sequences  $\{U_n(\alpha), n = 1, 2, \dots\}$

be  $\phi$ -mixing uniformly over all  $\alpha$ . This assumption guarantees that the series in (3.2) converges and

$$\lim_{n \rightarrow \infty} \frac{1}{n} E \left[ \left( \sum_{k=1}^n (U_k(\alpha) - v(\alpha)) \right)^2 \right] = \sigma^2(\alpha).$$

Assumption (3.5) concerns the convergence of this normalized variance term as  $\alpha \rightarrow \infty$ .

Put

$$Z_\alpha(t) = d(\alpha) \sigma(\alpha)^{-1} \sum_{k=1}^{[n(\alpha)t]} U_k(\alpha), \quad t \geq 0$$

where  $[x]$  is the largest integer less than or equal to  $x$ .

Let  $\mathcal{D}([0, \infty))$  be the collection of real-valued, right continuous functions on  $[0, \infty)$  which have left limits everywhere endowed with the metric given in Lindvall [8].

(3.9) THEOREM. If (3.4)-(3.6) are satisfied, then the process  $Z_\alpha = \{Z_\alpha(t); t \geq 0\}$  converges weakly in  $\mathcal{D}([0, \infty))$  to a Brownian motion with negative unit drift.

The proof of this result will be given in the next section.

Let

$$W(\alpha) = \sup_{n \geq 0} \sum_{k=1}^n U_k(\alpha).$$

Then it follows from Theorem (3.9) that



$$\lim_{\alpha \rightarrow \infty} P \left\{ \frac{d(\alpha)}{\sigma(\alpha)} W(\alpha) \leq x \right\} = \lim_{\alpha \rightarrow \infty} P \left\{ \sup_{t \geq 0} Z_{\alpha}(t) \leq x \right\}$$

$$= 1 - \exp\{-2x\}$$

where the last equality follows from the continuous mapping theorem (cf. Billingsley [1]). Thus we have the following result.

(3.10) THEOREM. If (3.4)-(3.6) hold, then

$$\lim_{\alpha \rightarrow \infty} P\{|v(\alpha)| W(\alpha) \leq x\} = 1 - \exp\{-2\sigma^{-2}x\}.$$

Hence, if the traffic intensity  $\lambda\mu^{-1}$  of a queue is less than but close to 1 and  $\{U_n\}$  is a stationary sequence satisfying (3.7) and (3.8), then the distribution of  $W$  is approximately exponential with mean  $-\frac{1}{2}\sigma^2 E[U_1]^{-1}$  where

$$\sigma^2 = \text{Var}[U_1] + 2 \sum_{n=1}^{\infty} \text{Cov}(U_1, U_{1+n}).$$

Thus positive cross correlation between the interarrival and service times will tend to decrease the average mean waiting time; while positive correlation within the interarrival time or service time sequence will tend to increase the mean waiting time.

(2.11) EXAMPLE. For each  $\alpha > 0$ , let  $Q_\alpha$  be a queue as in Example (2.1) with  $\{X_n(\alpha)\}$  (respectively  $\{S_n(\alpha)\}$ ) forming an EARMA(1,1) process with parameters  $\lambda(\alpha)$ ,  $\beta_1$  and  $\rho_1$  (respectively  $\mu(\alpha)$ ,  $\beta_2$  and  $\rho_2$ ). Assume  $\lambda(\alpha) < \mu(\alpha)$  for each  $\alpha$ , but  $\lim_{\alpha \rightarrow \infty} \lambda(\alpha) = \lim_{\alpha \rightarrow \infty} \mu(\alpha) = m$ ; then hypotheses (3.4)-(3.6) are satisfied and Theorem (3.10) holds with

$$\sigma^2 = 2m^{-2} \{ 1 + (1-\rho_1)^{-1} (1-\beta_1) [(1-\rho_1)\beta_1 + (1-\beta_1)\rho_1] + (1-\rho_2)^{-1} (1-\beta_2) [(1-\rho_2)\beta_2 + (1-\beta_2)\rho_2] \}.$$

Hence, if the traffic intensity  $\lambda\mu^{-1}$  of a queue with independent EARMA(1,1) services and interarrival times is less than but close to 1, then the distribution of the equilibrium waiting time  $W$  is approximately exponential with mean  $-\frac{1}{2} \sigma^2 E[U_1]$  where  $E[U_1] = \mu^{-1} - \lambda^{-1}$  and  $\sigma^2$  is given by

$$\sigma^2 = \lambda^{-2} \{ 1 + 2(1-\rho_1)^{-1} (1-\beta_1) [(1-\rho_1)\beta_1 + (1-\beta_1)\rho_2] \} + \mu^{-2} \{ 1 + 2(1-\rho_2)^{-1} (1-\beta_2) [(1-\rho_2)\beta_2 + (1-\beta_2)\rho_2] \}.$$

Thus the dependence in this queue tends to increase the mean customer waiting time at least in close to heavy traffic conditions.



(3.12) EXAMPLE. For each  $\alpha > 0$ , let  $Q_\alpha$  be a queue as in Example (2.7) with parameters  $E[X_n(\alpha)] = \lambda(\alpha)^{-1}$ ,  $\beta$ , and  $\rho$  and  $E[S_n(\alpha)] = \mu(\alpha)^{-1}$ . Again we will assume  $\lambda(\alpha) < \mu(\alpha)$  for each  $\alpha$  but  $\lim_{\alpha \rightarrow \infty} \lambda(\alpha) = \lambda = \lim_{\alpha \rightarrow \infty} \mu(\alpha) = \mu = m$ . Hence (3.4)-(3.6) are satisfied and Theorem (3.10) holds with

$$\begin{aligned}\sigma^2 &= \lambda^{-2} + \mu^{-2} - 2(1-\beta)(1-\rho)(\lambda\mu)^{-1} \\ &\quad + 2\rho(1-\rho)^{-1} [(1-\beta)^2\mu^{-2} - (1-\beta)(1-\rho)(\lambda\mu)^{-1}] \\ &= \lambda^{-2} + \mu^{-2} [1 + 2(1-\beta)^2\rho(1-\rho)^{-1}] - 2(\lambda\mu)^{-1}(1-\beta) \\ &= 2m^{-2} [1 + (1-\beta)^2\rho(1-\rho)^{-1} - (1-\beta)] .\end{aligned}$$

The quantity  $\sigma^2 - \sigma_{\text{ind}}^2$  can be either positive or negative where  $\sigma_{\text{ind}}^2 = \lambda^{-2} + \mu^{-2} = 2m^{-2}$ . It is negative if  $\rho < (2-\beta)^{-1}$  and nonnegative otherwise. Hence, if  $\rho$  is sufficiently less than  $(2-\beta)^{-1}$ , then the average customer waiting time in the queue will tend to be less than the waiting time in the independent case at least in close to heavy traffic conditions. It will tend to be greater if  $\rho$  is sufficiently greater than  $(2-\beta)^{-1}$ .

(3.13) EXAMPLE. For each  $\alpha > 0$ , let  $Q_\alpha$  be a queue as in Example (2.13) with parameters  $\lambda(\alpha)$ ,  $\mu(\alpha)$ ,  $\rho$ ,  $\beta$ , and  $\gamma$ . Assume  $\lim_{\alpha \rightarrow \infty} \lambda(\alpha) = \lambda = \lim_{\alpha \rightarrow \infty} \mu(\alpha) = \mu = m$  as before. Then (3.4)-(3.6) are satisfied and Theorem (3.10) holds with



$$\begin{aligned}
 \sigma^2 &= \mu^{-2} + \lambda^{-2} - 2(\lambda\mu)^{-1} (1-\beta)(1-\gamma) \\
 &\quad + 2\rho(1-\rho)^{-1} \{\mu^{-1}(1-\gamma) - \lambda^{-1}(1-\beta)\}^2 \\
 &= 2m^{-2} [1 - (1-\beta)(1-\gamma) + \rho(1-\rho)^{-1} \{(\beta-\gamma)\}^2] .
 \end{aligned}$$

The difference  $\sigma^2 - \sigma_{\text{ind}}^2$  can be either positive or negative; ( $\sigma_{\text{ind}}^2 = \lambda^{-2} + \mu^{-2} = 2m^{-2}$ ). The difference will be negative if  $\rho < (1-\beta)(1-\gamma)/[(1-\beta)(1-\gamma) + (\beta-\gamma)^2]$  and nonnegative otherwise. Hence under close to heavy traffic conditions, the average customer waiting time will tend to be less than the independent case if  $(1-\beta)(1-\gamma)/[(1-\beta)(1-\gamma) + (\beta-\gamma)^2]$  is sufficiently greater than  $\rho$ . It will be greater than the independent case if the same quantity is sufficiently less than  $\rho$ .

#### 4. PROOF OF THE MAIN RESULT

In this section we will give a proof of Theorem (3.9). We will assume that (3.4)-(3.6) are satisfied. Let

$$Y_\alpha(t) = \frac{d(\alpha)}{\sigma(\alpha)} \sum_{k=1}^{[n(\alpha)t]} [U_k(\alpha) - v(\alpha)] .$$

By Theorem 3' of Lindvall [8] and the continuous mapping theorem, Theorem (3.9) will be proved if for each  $a > 0$  we can show the following result.

(4.1) THEOREM. The process  $\{Y_\alpha(t); 0 \leq t \leq a\}$  converges weakly on  $\mathcal{D}([0,a])$  to Brownian motion.

( $\mathcal{D}([0,a])$  is the space of real-valued right continuous functions on  $[0,a]$  with left-hand limits endowed with the Skorohod  $J_1$  topology.)

We will prove Theorem (4.1) for  $a = 1$  by a series of lemmas. The proof for arbitrary  $a > 0$  is similar.

(4.2) LEMMA. The process  $\{Y_\alpha(t); 0 \leq t \leq 1\}$  has asymptotically independent increments.

Proof. Let  $0 \leq s_1 \leq t_1 < s_2 \leq t_2 < \dots < s_r \leq t_r \leq 1$ . Let  $C_i$  be the event  $\{Y_\alpha(t_i) - Y_\alpha(s_i) \leq x_i\}$  for real  $x_i$ . Then  $C_i$  is an event in the  $\sigma$ -algebra generated by

$U_{[n(\alpha)s_i]+1}, \dots, U_{[n(\alpha)t_i]}$ . If  $\delta$  is the smallest difference  $s_i - t_{i-1}$ , then  $[n(\alpha)s_i] + 1 - [n(\alpha)t_{i-1}] \geq [n(\alpha)\delta]$ . By (3.7)

$$|P(\bigcap_{i=1}^r C_i) - \prod_{i=1}^r P(C_i)| \leq r\phi([n(\alpha)\delta]).$$

Since  $\delta > 0$ ,  $r\phi([n(\alpha)\delta])$  tends to zero as  $\alpha \rightarrow \infty$  and the result follows.

In the following results, let  $D_i(\alpha) = U_i(\alpha) - v(\alpha)$  and  $V_m(\alpha) = \sum_{i=1}^m D_i(\alpha)$ . By (3.4)-(3.6) the following result holds.

(4.3) LEMMA.

$$a) \quad \frac{1}{K} E[V_k(\alpha)^2] \leq 2 \sum_{j=0}^{\infty} |E[D_1(\alpha) D_{1+j}(\alpha)]|;$$

$$b) \quad \lim_{\alpha \rightarrow \infty} n(\alpha)^{-1} E[V_{n(\alpha)}(\alpha)^2] = \sigma^2;$$

$$c) \quad \text{If } D_1 \text{ is bounded by } C, \text{ then } E[V_{n(\alpha)}(\alpha)^4] \leq K_\phi C^4 n(\alpha)^2$$

where  $K_\phi$  depends on  $\phi$  alone.

(4.4) LEMMA. The sequence of random variables  $\{n(\alpha)^{-1}(V_{n(\alpha)}(\alpha))^2; \alpha \geq 0\}$  is uniformly integrable.

Proof. By (3.4),  $\{D_1^2(\alpha); \alpha \geq 0\}$  is uniformly integrable. The remainder of the proof is very similar to that of (20.48) on page 176 of Billingsley [1] and will be omitted.



(4.5) LEMMA. For each positive  $\varepsilon$  there exists a  $\gamma$ ,  $\gamma > \sigma$  and an integer  $n_0$  such that  $n(\alpha) \geq n_0$  implies

$$P\left\{\max_{i \leq n(\alpha)} |V_i(\alpha)| \geq \gamma \sqrt{n(\alpha)}\right\} \leq \frac{\varepsilon}{\gamma^2}.$$

Proof. Put  $C_i^*(\alpha) = \sum_{j=1}^i |D_j(\alpha)|$ . By (3.4), there exists an increasing sequence of integers  $m_i$  such that  $n(\alpha) \geq m_i$  implies  $P\{|D_1(\alpha)| \geq \gamma \sqrt{n(\alpha)}/i^2\} \leq (\gamma n(\alpha) i^2)^{-1}$  for each positive  $\gamma$ . Let  $p(\alpha) = i$  if  $m_i \leq n(\alpha) < m_{i+1}$ ; then  $p(\alpha)$  tends to infinity as  $\alpha \rightarrow \infty$  but

$$(4.6) \quad \lim_{\alpha \rightarrow \infty} n(\alpha) P\{C_{p(\alpha)}^*(\alpha) \geq \gamma \sqrt{n(\alpha)}\}$$

$$\leq \lim_{\alpha \rightarrow \infty} n(\alpha) p(\alpha) P\{|D_1(\alpha)| \geq \gamma \sqrt{n(\alpha)} p(\alpha)^{-1}\}$$

$$\leq \lim_{\alpha \rightarrow \infty} n(\alpha) p(\alpha) P\{|D_1(\alpha)| \geq \gamma \sqrt{n(\alpha)} p(\alpha)^{-2}\}$$

$$\leq \lim_{\alpha \rightarrow \infty} n(\alpha) p(\alpha) [\gamma n(\alpha) p(\alpha)^2]^{-1} = 0.$$

We can further choose  $p(\alpha)$  so that  $p(\alpha) \leq n(\alpha)$ . Fix  $\varepsilon > 0$ .

By the previous lemma we can choose  $\gamma$  so that  $\gamma > \sigma$  and

$$P\{ \max_{i \leq n(\alpha)} |V_i(\alpha)| \geq 3\gamma \sqrt{n(\alpha)} \}$$

$$\leq P\{|V_{n(\alpha)}(\alpha)| \geq \gamma \sqrt{n(\alpha)}\}$$

$$+ [n(\alpha) + p(\alpha)] P\{C_{p(\alpha)}^*(\alpha) \geq \gamma \sqrt{n(\alpha)}\}$$

$$+ \sum_{i=1}^{n(\alpha)-p(\alpha)-1} P(\Lambda_i(\alpha)) [P\{|V_{n(\alpha)}(\alpha) - V_{i+p(\alpha)}(\alpha)| \geq \gamma \sqrt{n(\alpha)}\} + \phi(p(\alpha))]$$

$$\leq 2\epsilon\gamma^{-2} + \phi(p(\alpha)) + [n(\alpha) + p(\alpha)] P\{C_{p(\alpha)}^*(\alpha) \geq \gamma \sqrt{n(\alpha)}\}$$

by (4.7) and the fact that the  $\Lambda_i(\alpha)$  are disjoint. Hence by (4.6) and (3.8) for  $\alpha$  large

$$P\{ \max_{i \leq n(\alpha)} |V_i(\alpha)| \geq 3\gamma \sqrt{n(\alpha)} \} \leq 3\epsilon\gamma^{-2}.$$

PROOF OF THEOREM (4.1). For  $a = 1$ : By Lemma (4.5) the sequence  $\{Y_\alpha\}$  is tight on  $\mathbb{D}[0,1]$ . Further  $\lim_{\alpha \rightarrow \infty} E[Y_\alpha(t)] = 0$  and

$$\lim_{\alpha \rightarrow \infty} E[Y_\alpha(t)^2] = \lim_{\alpha \rightarrow \infty} (\sqrt{n(\alpha)} \sigma(\alpha))^{-2} E \left[ \left( \sum_{k=1}^{[n(\alpha)t]} D_k(\alpha) \right)^2 \right] = t$$

by (3.5). Since  $Y_\alpha(t)^2 = [\sigma^2(\alpha) n(\alpha)]^{-1} V_{[n(\alpha)t]}(\alpha)^2$ ,  $\{Y_\alpha^2(t)\}$  is uniformly integrable for each  $t$  by Lemma (4.4). By Lemma (4.2)  $\{Y_\alpha(t); 0 \leq t \leq 1\}$  has asymptotically independent increments. The result now follows from Theorem (19.2) of Billingsley [1].

$$(4.7) \quad P\{|V_{k(\alpha)}(\alpha)| > \gamma \sqrt{k(\alpha)}\} < \epsilon \gamma^{-2}$$

for all  $\alpha$ . Let

$$\Lambda_i(\alpha) = \{\max_{j < i} |V_j(\alpha)| < 3\gamma \sqrt{n(\alpha)} \leq |V_i(\alpha)|\};$$

then

$$P\{\max_{i \leq n(\alpha)} |V_i(\alpha)| \geq 3\gamma \sqrt{n(\alpha)}\}$$

$$\leq P\{|V_{n(\alpha)}(\alpha)| \geq \gamma \sqrt{n(\alpha)}\}$$

$$+ \sum_{i=1}^{n(\alpha)-1} P(\Lambda_i(\alpha) \cap \{|V_{n(\alpha)}(\alpha) - V_i(\alpha)| \geq 2\gamma \sqrt{n(\alpha)}\}).$$

The summation is bounded above by

$$\sum_{i=1}^{n(\alpha)-p(\alpha)-1} P\{|V_i(\alpha) - V_{i+p(\alpha)}(\alpha)| \geq \gamma \sqrt{n(\alpha)}\}$$

$$+ \sum_{i=1}^{n(\alpha)-p(\alpha)-1} P(\Lambda_i(\alpha) \cap \{|V_{n(\alpha)}(\alpha) - V_{i+p(\alpha)}(\alpha)| > \gamma \sqrt{n(\alpha)}\})$$

$$+ \sum_{i=n(\alpha)-p(\alpha)}^{n(\alpha)-1} P\{|V_{n(\alpha)}(\alpha) - V_i(\alpha)| \geq \gamma \sqrt{n(\alpha)}\}.$$

Each term in the first and third of these sums is bounded above by  $P\{C_{p(\alpha)}^*(\alpha) \geq \gamma \sqrt{n(\alpha)}\}$ . Since  $\Lambda_i(\alpha)$  is measurable with respect to the  $\sigma$ -algebra generated by  $U_1(\alpha), \dots, U_i(\alpha)$  by (3.7)



## 5. EXTENSIONS

This paper presented a heavy traffic result for the limiting distribution of the customer waiting time in single server queues with dependent interarrival and service times. There are many different ways in which to model dependent interarrival and service times in queues using the EARMA scheme. Some other ways are given in Lewis and Shedler [7]. Properties of these multivariate exponential models will be given elsewhere.

Work is currently going on in the simulation of these queues. Some simulation results have been reported for the queue of Example (2.7) with  $\beta = 0.5$ ,  $\mu = 4$  and various values of  $\rho$  and  $\lambda$  in Boonsong [2]. They seem to indicate that the heavy traffic result approximates the average customer waiting time fairly well for a traffic intensity of .9 and quite well for a traffic intensity of .95. However, the distribution of  $W$  in a simulation with  $\rho = 0.25$  and traffic intensity .995 shows a large amount of underdispersion relative to an exponential distribution. Thus, it seems that the convergence of the waiting time distribution is very slow.

It is expected that further simulations will give more indication of the validity of using the heavy traffic result to approximate the average customer waiting time, or quantiles of the waiting time distribution under less than heavy traffic conditions. These results will be reported on elsewhere.

## 6. REFERENCES

- [1] BILLINGSLEY, P. (1968). Convergence of Probability Measures. Wiley, New York.
- [2] BOONSONG, P. (1978). A Simulation Study of a Class of First-Come First-Served Queues with EARMA Correlation Structure, Master Thesis, Naval Postgraduate School, Monterey, California, September.
- [3] JACOBS, P. A. and LEWIS, P.A.W. (1977). A mixed autoregressive-moving average exponential sequence and point process (EARMA 1,1). Adv. Appl. Prob. 9, 87-104.
- [4] JACOBS, P. A. (1978). A cyclic queueing network with dependent exponential service times. J. Appl. Prob. 15, 573-589.
- [5] KINGMAN, J. F. C. (1962). On queues in heavy traffic. J. Roy. Statist. B 24, 383-392.
- [6] LEWIS, P. A. W. and SHEDLER, G. S. (1977). Analysis and modelling of point processes in computer systems. Bull. Int. Statist. Inst. 47, 193-210.
- [7] LEWIS, P.A.W. and SHEDLER, G.S. (1977). Analysis and modelling of point processes in computer systems. Naval Postgraduate School Technical Report NPS55-77-38.
- [8] LINDVALL, T. (1973). Weak convergence of probability measures and random functions in the function space  $D[0, \infty)$ . J. Appl. Prob. 10, 109-121.
- [9] LOYNES, R.M. (1962). The stability of a queue with non-independent interarrival and services. Proc. Camb. Philos. Soc. 58, 497-520.
- [10] ROSENBLATT, M. (1971). Markov Processes. Structure and Asymptotic Behavior. Springer-Verlag, Berlin.



# DISTRIBUTION LIST

Defense Documentation Center  
Cameron Station  
Alexandria, VA 22314

Technical Information Division  
Naval Research Laboratory  
Washington, D.C. 20375

Library Code 0212  
Naval Postgraduate School  
Monterey, CA 93940

Library Code 55  
Naval Postgraduate School  
Monterey, CA 93940

Dean of Research  
Code 012  
Naval Postgraduate School  
Monterey, CA 93940

Statistics and Probability Program  
Office of Naval Research  
Code 436  
Arlington, VA 22217

Library  
Naval Ocean Systems Ctr  
San Diego, CA 92152

Naval Electronic Systems Command  
Navelex 320  
National Center No. 1  
Arlington, VA 20360

Director Naval Research Lab  
Attn: Library (ONRL)  
Code 2029  
Washington, D.C. 20375

Office of Naval Research  
San Francisco Area Office  
760 Market Street  
San Francisco, CA 94102

No. of Copies

2

1

2

1

1

1

1

1

1

1



# No. of Copies

Prof. T. W. Anderson  
Department of Statistics  
Stanford University  
Stanford, CA 94305

1

Professor F. J. Anscombe  
Department of Statistics  
Yale University  
New Haven, CT 06520

1

Professor C. R. Baker  
Department of Statistics  
University of North Carolina  
Chapel Hill, N.C. 37512

1

Prof. R. E. Barlow  
Operations Research Ctr  
College of Engineering  
University of California  
Berkeley, CA 94720

1

Prof. R. E. Bechhofer  
Department of Operations Research  
Cornell University  
Ithaca, New York 14850

1

Mr. C. N. Bennett  
Naval Coastal Systems Lab  
Code P761  
Panama City, FL 32401

1

Prof. N. J. Bershad  
School of Engineering  
University of California  
Irvine, CA 92664

1

Prof. L. N. Bhat  
Comput Sci/Oper Res Ctr  
Southern Methodist Univ.  
Dallas, TX 75275

1

P. J. Bickel  
Department of Statistics  
University of California  
Berkeley, CA 94720

1

Prof. W. R. Blischke  
Dept. of Quantitative Business Analysis  
Univ. of Southern California  
Los Angeles, CA 90007

1

No. of Copies

No. of Copies

Prof. H. W. Block  
Dept. of Mathematics  
University of Pittsburgh  
Pittsburgh, PA 15260

1

Prof. Peter Bloomfield  
Dept. of Statistics  
Princeton University  
Princeton, N.J. 08540

1

Prof. Joseph Blum  
Dept. of Math., Stat. and Com. Sci.  
The American University  
Washington, D.C. 20016

1

J. E. Boyer, Jr.  
Dept. of Statistics  
Southern Methodist University  
Dallas, TX 75275

1

Dr. William Brogan  
National Science Foundation  
Systems Theory and Applications Program Eng. Div.  
18th and "G", N.W.  
Washington, D.C. 20550

1

Prof. Ted Brown  
Dept. of Computer Science  
Queens College  
Flushing, N.Y. 11367

1

Dr. J. Chandra  
U. S. Army Research  
P.O. Box 12211  
Research Triangle Park  
North Carolina 27706

1

Prof. H. Chernoff  
Dept. of Mathematics  
Mass. Inst. Tech.  
Cambridge, MA 02139

1

Prof. E. Çinlar  
Dept. of Industrial Engineering  
Northwestern University  
Evanston, ILL 60201

1

Dr. D. R. Cox  
Dept. of Mathematics  
Imperial College  
London SW7 2BZ, ENGLAND

1



No. of Copies

No. of Copies

Prof. C. Derman  
Dept. of Civil Eng. and Eng. Mech.  
Columbia University  
New York, New York 10027

Prof. R. L. Disney  
Virginia Poly. Inst. and State Univ.  
Dept. of Ind. Eng.  
Blacksburg, VA 24061

Mr. J. Dowling  
Defense Logistics Studies  
Information Exchange  
Army Logistics Management Center  
Fort Lee, VA 20390

Dr. M. J. Fischer  
Defense Communications Agency  
1860 Wiehle Ave.  
Reston, VA 22070

Prof. George S. Fishman  
Curr. in OR and Sys Anal.  
Univ. of North Carolina  
Phillips Annex  
Chapel Hill, N.C. 20742

Prof. Larry George  
Ind. Eng. Dept.  
Texas A&M University  
College Station, TX 77843

Mr. Gene H. Gleissner  
Appl. Math. Lab.  
David Taylor Naval Ship Research  
and Devel. Ctr  
Bethesda, MD 20084

Dr. R. Gnanadesikan  
Bell Telephone Lab.  
Murray Hill, N.J.

Dr. A. J. Goldman, Chief OR  
Div. 205.02, Admin. A428  
U.S. Dept. of Commerce  
Washington, D.C. 20234

Prof. S. S. Gupta  
Dept. of Statistics  
Purdue University  
Lafayette, IN 47907



No. of Copies

Prof. D. L. Hanson  
Dept. of Math. Sci.  
State University of New York,  
Binghamton, NY 13901

1

Prof. F. J. Harris  
Dept. of Elec. Eng.  
San Diego State Univ.  
San Diego, CA 92182

1

Prof. L. H. Herbach  
Dept. of Oper. Res. and Sys. Anal.  
Polytechnic Inst. of N.Y.  
Brooklyn, NY 11201

1

Prof. M. J. Hinich  
Dept. of Economics  
Virginia Polytechnic Inst. and State Univ.  
Blacksburg, VA 24061

1

Prof. W. M. Hirsch  
Inst. of Math. Sci.  
New York Univ.  
New York, N.Y. 19453

1

Prof. D. L. Iglehart  
Dept. of Oper. Res.  
Stanford Univ.  
Stanford, CA 94350

1

Prof. J. B. Kadane  
Dept. of Statistics  
Carnegie-Mellon  
Pittsburgh, PA 15213

1

Dr. H. Kobayashi  
IBM  
Yorktown Heights, N.Y. 10598

1

Dr. Guy Latouche  
Laboratoire d'Informatique Theorique  
Universite Libre de Bruxelles  
Boulevard du Triomphe  
1050 Bruxelles, BELGIUM

1

Dr. Richard Lau, Director  
Office of Naval Research Branch Office  
1030 Green Street  
Pasadena, CA 91101

1

1

1

1

1

1

1

2

1

1

1



Prof. J. A. Muckstadt  
Dept. of Oper. Res.  
Cornell University  
Ithaca, N.Y. 19850

Dr. Janet M. Myhre  
Inst. Dec. Sci. for Bus. and Pub. Pol.  
Claremont Men's College  
Claremont, CA 91711

Prof. Marcel Neuts  
University of Delaware  
Dept. of Mathematics  
Newark, Delaware 19711

W. R. Nunn  
Center for Naval Analyses  
1401 Wilson Blvd  
Arlington, VA 22209

B. S. Orleans  
Naval Sea Systems Command  
(Sea 03F), Rm 10SC8  
Arlington, VA 20360

Prof. D. B. Owen  
Dept. of Statistics  
Southern Methodist Univ.  
Dallas, TX 75222

Prof. B. Pagurek  
Dept. of Sys Eng and Com Sci  
Carleton University  
Ottawa, Ont K1S 5B6  
Canada

Prof. E. Parzen  
Statistics Dept.  
Texas A&M  
College Station, TX 77843

Prof. S. L. Phoenix  
Sibley School of Mech. and Aero. Eng.  
Cornell University  
Ithaca, N.Y. 14850

Prof. M. L. Puri  
Dept. of Mathematics  
P.O. Box F  
Indiana Univ. Foundation  
Bloomington, IN 47401



# No. of Copies

Prof. M. Rosenblatt  
Dept. of Mathematics  
Univ. of California  
San Diego, CA 92093

1

Prof. S. M. Ross  
College of Engineering  
Univ. of California  
Berkeley, CA 94720

1

M. N. Rothkopf  
Xerox Palo Alto Research Ctr  
333 Coyote Hill Road  
Palo Alto, CA 94304

1

Robert Sargent  
Dept. of Ind. Eng. and Oper. Res.  
Syracuse Univ.  
Syracuse, N.Y. 13210

1

Prof. L. L. Scharf Jr.  
Dept. of Elec. Eng.  
Colorado State University  
Ft. Collins, CO 89521

1

Dr. Paul Schwitzer  
Graduate School of Management  
Univ. of Rochester  
Rochester, N.Y. 14627

1

Prof. D. G. Shimshak  
Univ. of Massachusetts  
Harbor Campus  
Boston, MA 02125

1

Prof. D. C. Siegmund  
Dept. of Statistics  
Stanford Univ.  
Stanford, CA 94305

1

Prof. W. L. Smith  
Dept. of Statistics  
Univ. of North Carolina  
Chapel Hill, N.C. 27514

1

Prof. J. G. Shanthikumar  
Dept. of Ind. Eng.  
University of Toronto  
Toronto, Ont. M5S 1R4  
CANADA

1

|   | No. of Copies |
|---|---------------|
| Prof. J. R. Thompson<br>Dept. of Math. Sci.<br>Rice University<br>Houston, TX 77001             | 1             |
| Prof. J. L. Tukey<br>Dept. of Statistics<br>Princeton Univ.<br>Princeton, N.J. 98540            | 1             |
| Daniel H. Wagner<br>Station Square One<br>Paoli, PA 19301                                       | 1             |
| Prof. Grace Wahba<br>Dept. of Statistics<br>Univ. of Wisconsin<br>Madison, WI 53706             | 1             |
| Prof. G. S. Watson<br>Dept. of Statistics<br>Princeton Univ.<br>Princeton, N.J. 08540           | 1             |
| Prof. Roy Welsch<br>M.I.T.<br>Sloan School of Management<br>Cambridge, MA 02139                 | 1             |
| Naval Postgraduate School<br>Monterey, CA. 93940<br>Attn: Prof. D.P. Gaver, Code 55Gv           | 1             |
| Prof. P.A. Jacobs, Code 55Jc  | 20            |
| R. J. Stampfel, Code 55   | 1             |
| Prof. R. A. Bradley<br>Dept. of Statistics<br>Florida State University<br>Tallahassee, FL 32306 | 1             |